

Generalization of Some Arithmetical Properties of Fermat-Euler Dynamical Systems

Ahmed N. Elsaywy*

October 29, 2009

Abstract

We study and generalize some arithmetical properties of the classes (2^k+) and (2^k-) introduced by V. I. Arnold: a number n belongs to the class $(N+)$ if $N|\varphi(n)$ and $2^{\frac{\varphi(n)}{N}} \equiv 1 \pmod{n}$ where $\varphi(n)$ is the Euler function, and belongs to the class $(M-)$ if $M|\varphi(n)$ and $2^{\frac{\varphi(n)}{M}} \equiv -1 \pmod{n}$. The classes $(2\pm)$, $(4\pm)$ and $(8\pm)$ are studied by V. I. Arnold in [1, 2] and here we will show general properties of the classes $(2^k\pm)$ and we will see that the properties which proved by V. I. Arnold are special cases of ours.

1 Introduction

One of the most important theorems in number theory and theory of finite groups is Fermat's little theorem ($a^{p-1} \equiv 1 \pmod{p}$, where p is prime and $a \not\equiv 0 \pmod{p}$ is an integer) and its Euler's generalization ($a^{\varphi(n)} \equiv 1 \pmod{n}$, where a and n are coprime integers). They are also some of the most important applications of the dynamical system theory. The Fermat-Euler dynamical system is an example of the connection between dynamical systems and number theory in which we try to use the properties of the dynamical systems to generalize or get new properties in number theory.

Definition: The Abelian multiplicative group of residues coprime to n is called the *Euler group* $\Gamma(n)$, and the *Euler function* $\varphi(n)$ is its order.

The Euler function is calculated as follows:

$$\varphi(1) = 1 \text{ and } \varphi(n) = \prod (p_i - 1)p_i^{a_i-1},$$

*Adresse: Mathematisches Institut, Heinrich-Heine-Universität, D-40225 Düsseldorf, Germany, Email: elsaywy@math.uni-duesseldorf.de

where $n = \prod p_i^{a_i}$ is the prime factorization of n , and has the property $\varphi(mn)\varphi(d) = \varphi(m)\varphi(n)d$ where $d = (m, n)$.

Definition: A Fermat-Euler dynamical system is a dynamical system with the ring of residues modulo n as its set and the function $f(x) = ax$, where $(a, n) = 1$, as its function.

In what follows, we consider the case in which $a = 2$ and n is odd. In [1], V. I. Arnold has proved the following theorem which is a generalization of Euler's Theorem, by using the Fermat-Euler dynamical system.

Theorem A[1]. All cycles of the Fermat-Euler permutation have the same length (period) T . Therefore, the period and the number $N = N(n)$ of cycles satisfy the relation

$$\varphi(n) = NT \quad \text{and} \quad a^{\varphi(n)/N} \equiv 1 \pmod{n}.$$

This theorem was the starting point of defining and studying new classes of numbers called $(N\pm)$.

Definition: An odd number n belongs to the class $(N+)$ if $N|\varphi(n)$ and

$$2^{\varphi(n)/N} \equiv +1 \pmod{n},$$

and belongs to the class $(M-)$ if $M|\varphi(n)$ and

$$2^{\varphi(n)/M} \equiv -1 \pmod{n}.$$

The following properties directly follow from the definition

1. If $K | N$, then $(N+) \subseteq (K+)$.
2. If M is odd, then $(M-)$ is empty.
3. If $M = K(2m+1)$, then $(M-) \subseteq (K-)$.
4. $(2N-) \subseteq (N+)$.

The proof of these properties can be found in [1].

In [1], V. I. Arnold calculated for each odd integer $1 < n < 512$ its classes $(N+)$ and $(M-)$ with maximal values of N and M as well as the values of its minimal period $T(n)$ and the Euler function $\varphi(n)$. From these calculations he observed and proved some properties of the classes $(2\pm)$, $(4\pm)$ and $(8\pm)$. In this work we give and prove general properties of the classes (2^k+) and

(2^k-) , where k is a positive integer, and show that the properties which are proved in [1, 2] by V. I. Arnold are special cases of the generalized properties.

In [2], V. I. Arnold proved by induction that if n is divisible by more than k distinct primes, then $n \in (2^k+)$. We will give here another proof and we will answer the following question.

Question: When does n belong to (2^k+) or (2^k-) if n has exactly k or $k-1$ distinct prime divisors, and when does it not?

In section 2, we prove the following theorems which give the complete answer to this question.

Theorem 1. Let n be an odd integer and have exactly $k \geq 2$ distinct prime divisors.

- If $n = \prod_{i=1}^k p_i^{a_i}$ with $p_i \equiv 3 \pmod{8}$ for all i , then $n \in (2^k-)$.
- If $n = \prod p_i^{a_i} \prod q_j^{b_j}$ with $p_i \equiv 3 \pmod{8}$ for all i , and $q_j \equiv -1 \pmod{8}$ for all j , then $n \notin (2^k+) \cup (2^k-)$.
- If $n = q^b \prod_{i=1}^{k-1} p_i^{a_i}$ with $p_i \equiv 3$ or $-1 \pmod{8}$, and $q \equiv -3 \pmod{8}$, then again $n \notin (2^k+) \cup (2^k-)$.
- If n is not in the previous three cases, then $n \in (2^k+)$.

Theorem 2. Let n be an odd integer and have exactly $k-1 \geq 2$ distinct prime divisors.

- If $k = 3$ and $n = p_1^{a_1} p_2^{a_2}$ with $p_1 \equiv p_2 \equiv -3 \pmod{8}$, then $n \in (2^k-)$.
- If $n = \prod_{i=1}^r p_i^{a_i} \prod_{j=r+1}^{k-1} q_j^{b_j}$ with $p_i \equiv -3 \pmod{8}$ for all i , $q_j \equiv -1$ or $3 \pmod{8}$ for all j , and $0 \leq r \leq 2$, then $n \notin (2^k+) \cup (2^k-)$.
- $n = p_1^{a_1} \prod_{j=2}^{k-1} q_j^{b_j}$ with $p_1 \in (4-)$ and $q_j \equiv -1$ or $3 \pmod{8}$ for all j , then $n \notin (2^k+) \cup (2^k-)$.
- If n is not in the previous three cases, then $n \in (2^k+)$.

In section 3 we will see the cases $k = 2$ and $k = 3$ as an example and how the theorems in section 2 describe the classes $(4\pm)$ and $(8\pm)$. Also we will compare between our results and the results which are proven by V. I. Arnold in [1, 2].

2 The classes $(2^k \pm)$

We will use the following propositions in our proofs, which can be found in any elementary book of Number Theory, for example in [3].

Proposition 1. *If $x \equiv a \pmod A$ and $x \equiv a \pmod B$, then $x \equiv a \pmod C$, where C is the least common multiple of A and B .*

Proposition 2. *Let p be an odd prime, then*

$$2^{\frac{\varphi(p^a)}{2}} \equiv \begin{cases} +1 \pmod{p^a} & \text{if } p \equiv \pm 1 \pmod 8 \\ -1 \pmod{p^a} & \text{if } p \equiv \pm 3 \pmod 8 \end{cases}$$

This proposition can be easily obtained from Euler's criterion ($a^{\frac{p-1}{2}} \equiv (\frac{a}{p}) \pmod p$, where $(\frac{a}{p})$ is the Legendre symbol). Indeed, since we know that

$$\left(\frac{2}{p}\right) \equiv \begin{cases} +1 \pmod p & \text{if } p \equiv \pm 1 \pmod 8 \\ -1 \pmod p & \text{if } p \equiv \pm 3 \pmod 8, \end{cases}$$

we can write $2^{\frac{p-1}{2}} = \pm 1 + Ap$ for some integer A . Hence, using binomial formula, we can easily get

$$2^{\frac{(p-1)p^{a-1}}{2}} = (\pm 1 + Ap)^{p^{a-1}} = \pm 1 + A'p^a, \text{ because } p^{a-1} \text{ is odd.}$$

The following theorem is proved in [2] by induction, we give here a direct proof using Proposition 2.

Theorem 2.1. *Every odd number which is divisible by more than k different primes belongs to the class $(2^k +)$.*

Proof. Let n be an odd number which is divisible by exactly r different primes with $0 < k < r$, then we can write n as $n = AB$ where $A = \prod_{i=1}^k p_i^{a_i}$ and $B = \prod_{i=k+1}^r p_i^{a_i}$.

From Euler's theorem, it follows that $2^{\varphi(B)} \equiv 1 \pmod B$. Also, note that $\varphi(A)/2^k$ is an integer, because A has k different (odd) prime divisors. So

$$2^{\varphi(n)/2^k} = 2^{\varphi(A)\varphi(B)/2^k} = (2^{\varphi(B)})^{\varphi(A)/2^k} \equiv 1 \pmod B.$$

Since $\varphi(B)$ is even, $\varphi(A) = \prod_{i=1}^k \varphi(p_i^{a_i})$, and $2^{\varphi(p_i^{a_i})/2} \equiv \pm 1 \pmod{p_i^{a_i}}$, we have

$$2^{\varphi(n)/2^k} = 2^{\varphi(A)\varphi(B)/2^k} = (2^{\varphi(p_i^{a_i})/2})^{\varphi(B) \frac{\prod_{j=1, j \neq i}^k \varphi(p_j^{a_j})}{2^{k-1}}} \equiv 1 \pmod{p_i^{a_i}}$$

for every $1 \leq i \leq k$.

Thus, using Proposition 1, we get $2^{\varphi(n)/2^k} \equiv 1 \pmod{\prod_{i=1}^k p_i^{a_i} = A}$.

Now we have $2^{\varphi(n)/2^k} \equiv 1 \pmod{A}$ and $2^{\varphi(n)/2^k} \equiv 1 \pmod{B}$ where A and B are relatively prime; therefore,

$$2^{\varphi(n)/2^k} \equiv +1 \pmod{AB = n}; \text{ that is, } n \in (2^k+). \quad \square$$

Theorem 2.1 and Proposition 2 give us, in particular, a complete description of the classes $(2\pm)$, i.e. the case $k = 1$, as follows.

- Every odd number which is divisible by more than one prime belongs to the class $(2+)$.
- If $n = p^a$ with $p \equiv \pm 1 \pmod{8}$, then $n \in (2+)$.
- If $n = p^a$ with $p \equiv \pm 3 \pmod{8}$, then $n \in (2-)$.

Now the question is, what will happen in the case when n has exactly k distinct prime divisors? In Theorems 2.2-2.6 we study this case and show precisely when n belongs to the class (2^k+) , to the class (2^k-) , and when n belongs to neither of them.

Theorem 2.2. *If $n = \prod_{i=1}^k p_i^{a_i}$ has at least one prime divisor congruent to 1 modulo 8, then $n \in (2^k+)$.*

Proof. WLOG, we assume that $p_1 \equiv 1 \pmod{8}$ and $n = p_1^{a_1} A$ where $A = \prod_{i=2}^k p_i^{a_i}$. Then

$$2^{\varphi(n)/2^k} = (2^{\frac{\varphi(p_1^{a_1})}{2}})^{\frac{\varphi(A)}{2^{k-1}}} \equiv (+1)^{\frac{\varphi(A)}{2^{k-1}}} \equiv 1 \pmod{p_1^{a_1}}.$$

For all $i = 2, \dots, k$, we have $2^{\varphi(p_i^{a_i})/2} \equiv +1$ or $-1 \pmod{p_i^{a_i}}$, and since $p_1 \equiv 1 \pmod{8}$, it follows that $\frac{p_1-1}{2}$ is even, so $\frac{\varphi(p_1^{a_1})}{2}$ is even. Thus,

$$2^{\varphi(n)/2^k} = (2^{\frac{\varphi(A)}{2^{k-1}}})^{\frac{\varphi(p_1^{a_1})}{2}} = (2^{\varphi(p_i^{a_i})/2})^{\frac{\varphi(p_1^{a_1})}{2} \frac{\prod_{j=2, j \neq i}^k \varphi(p_j^{a_j})}{2^{k-2}}} \equiv 1 \pmod{p_i^{a_i}}$$

for every $2 \leq i \leq k$.

It follows that $2^{\varphi(n)/2^k} \equiv +1 \pmod{A}$, by Proposition 1.

So we have $2^{\varphi(n)/2^k} \equiv 1 \pmod{p_1^{a_1}}$ and $2^{\varphi(n)/2^k} \equiv 1 \pmod{A}$ where $p_1^{a_1}$ and A are relatively prime. Thus,

$$2^{\varphi(n)/2^k} \equiv +1 \pmod{p_1^{a_1} A = n}; \text{ that is, } n \in (2^k+). \quad \square$$

Now assume that n has no prime divisors congruent to 1 modulo 8. In Theorems 2.3, and 2.4, we deal with the case when all the prime divisors of n have the same residue (except 1) modulo 8.

Theorem 2.3. *If $n = \prod_{i=1}^k p_i^{a_i}$ with either $p_i \equiv -1 \pmod{8}$ for all i or $p_i \equiv -3 \pmod{8}$ for all i , then $n \in (2^k+)$.*

Proof. Case 1. Let $p_i \equiv -1 \pmod{8}$ for all i , then by Proposition 2 we have $2^{\varphi(p_i^{a_i})/2} \equiv 1 \pmod{p_i^{a_i}}$ for all i , so

$$2^{\varphi(n)/2^k} = (2^{\varphi(p_i^{a_i})/2})^{\frac{\prod_{j=1, j \neq i}^k \varphi(p_j^{a_j})}{2^{k-1}}} \equiv (+1)^{\frac{\prod_{j=1, j \neq i}^k \varphi(p_j^{a_j})}{2^{k-1}}} \equiv 1 \pmod{p_i^{a_i}}.$$

Thus, $2^{\varphi(n)/2^k} \equiv +1 \pmod{n}$; that is, $n \in (2^k+)$.

Case 2. Let $p_i \equiv -3 \pmod{8}$ for all i , then $2^{\varphi(p_i^{a_i})/2} \equiv -1 \pmod{p_i^{a_i}}$, and $\varphi(p_i^{a_i})/2$ is even, for all i . Indeed, let $p_i = 5 + r_i 8$ for some integer r_i , then $\frac{p_i-1}{2} = 2 + r_i 4$ is even; therefore, $\varphi(p_i^{a_i})/2 = p_i^{a_i-1}(p_i-1)/2$ is even for all i .

Hence,

$$2^{\varphi(n)/2^k} = (2^{\varphi(p_i^{a_i})/2})^{\frac{\prod_{j=1, j \neq i}^k \varphi(p_j^{a_j})}{2^{k-1}}} \equiv (-1)^{\frac{\prod_{j=1, j \neq i}^k \varphi(p_j^{a_j})}{2^{k-1}}} \equiv +1 \pmod{p_i^{a_i}}.$$

So we have again $2^{\varphi(n)/2^k} \equiv +1 \pmod{n}$, or $n \in (2^k+)$. $\square$

Theorem 2.4. *If $n = \prod_{i=1}^k p_i^{a_i}$ with $p_i \equiv 3 \pmod{8}$ for all i , then $n \in (2^k-)$.*

Proof. Since $p_i \equiv 3 \pmod{8}$ for all i , it follows from Proposition 2 that $2^{\varphi(p_i^{a_i})/2} \equiv -1 \pmod{p_i^{a_i}}$, and $\varphi(p_i^{a_i})/2$ is odd, for all i . Indeed, let $p_i = 3 + r_i 8$ for some integer r_i , then $\frac{p_i-1}{2} = 1 + r_i 4$ is odd; therefore, $\varphi(p_i^{a_i})/2 = p_i^{a_i-1}(p_i-1)/2$ is odd for all i .

Thus,

$$2^{\varphi(n)/2^k} = (2^{\varphi(p_i^{a_i})/2})^{\frac{\prod_{j=1, j \neq i}^k \varphi(p_j^{a_j})}{2^{k-1}}} \equiv (-1)^{\frac{\prod_{j=1, j \neq i}^k \varphi(p_j^{a_j})}{2^{k-1}}} \equiv -1 \pmod{p_i^{a_i}}.$$

So $2^{\varphi(n)/2^k} \equiv -1 \pmod{\prod_{i=1}^k p_i^{a_i} = n}$; that is, $n \in (2^k-)$. $\square$

In the next two theorems, n has no prime divisor $p \equiv 1 \pmod{8}$ and has at least two prime divisors p and q with different residues modulo 8.

Theorem 2.5. *Let $n = \prod_{i=1}^r p_i^{a_i} \prod_{j=r+1}^k q_j^{b_j}$ with $0 < r < k$, $p_i \equiv 3$ or $-1 \pmod{8}$ for all i , and $q_j \equiv -3 \pmod{8}$ for all j , then*

$$\begin{aligned} n &\notin (2^k+) \cup (2^k-) && \text{if } r = k-1, \text{ and} \\ n &\in (2^k+) && \text{if } r < k-1. \end{aligned}$$

Proof. Note that $\varphi(p_i^{a_i})/2$ is odd. Indeed, $p_i \equiv 3$ or $-1 \pmod{8}$ means that $p_i \equiv 3 \pmod{4}$, so we can write $p_i = 3 + 4s_i$ for some integer s_i ; therefore, $\frac{p_i-1}{2} = 1 + 2s_i$, and $\varphi(p_i^{a_i})/2 = p_i^{a_i-1}(\frac{p_i-1}{2})$ is odd.

On the other hand, $\varphi(q_j^{b_j})/2$ is even. Indeed, $q_j \equiv -3 \pmod{8}$ means that $q_j \equiv 1 \pmod{4}$, and we can write $q_j = 1 + 4s_j$ for some integer s_j ; therefore, $\frac{q_j-1}{2} = 2s_j$, so $\varphi(q_j^{b_j})/2 = q_j^{b_j-1}(\frac{q_j-1}{2})$ is even.

So we have

$$\begin{aligned} 2^{\varphi(n)/2^k} &= (2^{\varphi(p_i^{a_i})/2})^{\frac{\prod_{l=1, l \neq i}^r \varphi(p_l^{a_l}) \prod_{j=r+1}^k \varphi(q_j^{b_j})}{2^{k-1}}} \equiv (\pm 1)^{\frac{\prod_{l=1, l \neq i}^r \varphi(p_l^{a_l}) \prod_{j=r+1}^k \varphi(q_j^{b_j})}{2^{k-1}}} \\ &\equiv +1 \pmod{p_i^{a_i}}, \text{ because } \varphi(q_j^{b_j})/2 \text{ is even.} \end{aligned}$$

Therefore, $2^{\varphi(n)/2^k} \equiv 1 \pmod{\prod_{i=1}^r p_i^{a_i}}$.

Case 1. Let $r = k - 1$; that is, $n = q^b \prod_{i=1}^{k-1} p_i^{a_i}$ with $p_i \equiv 3$ or $-1 \pmod{8}$, and $q \equiv -3 \pmod{8}$, then

$$2^{\varphi(n)/2^k} = (2^{\varphi(q^b)/2})^{\frac{\prod_{i=1}^{k-1} \varphi(p_i^{a_i})}{2^{k-1}}} \equiv (-1)^{\frac{\prod_{i=1}^{k-1} \varphi(p_i^{a_i})}{2^{k-1}}} \equiv -1 \pmod{q^b},$$

because $\frac{\prod_{i=1}^{k-1} \varphi(p_i^{a_i})}{2^{k-1}}$ is odd. Thus,

$$2^{\varphi(n)/2^k} \equiv +1 \pmod{\prod_{i=1}^r p_i^{a_i}} \text{ and } 2^{\varphi(n)/2^k} \equiv -1 \pmod{q^b}.$$

Now assume that $n \in (2^k+) \cup (2^k-)$, then $2^{\varphi(n)/2^k} \equiv \pm 1 \pmod{n}$; therefore, $2^{\varphi(n)/2^k}$ has the same residue modulo $\prod_{i=1}^r p_i^{a_i}$ and modulo q^b . But this contradicts our calculations, so $n \notin (2^k+) \cup (2^k-)$.

Case 2. Let $0 < r < k - 1$ and $n = \prod_{i=1}^r p_i^{a_i} \prod_{j=r+1}^k q_j^{b_j}$, then

$$\begin{aligned} 2^{\varphi(n)/2^k} &= (2^{\varphi(q_j^{b_j})/2})^{\frac{\prod_{i=1}^r \varphi(p_i^{a_i}) \prod_{l=r+1, l \neq j}^k \varphi(q_l^{b_l})}{2^{k-1}}} \equiv (-1)^{\frac{\prod_{i=1}^r \varphi(p_i^{a_i}) \prod_{l=r+1, l \neq j}^k \varphi(q_l^{b_l})}{2^{k-1}}} \\ &\equiv +1 \pmod{q_j^{b_j}}, \end{aligned}$$

because $\frac{\varphi(q_l^{b_l})}{2}$ is even. Thus,

$2^{\varphi(n)/2^k} \equiv 1 \pmod{\prod_{i=1}^r p_i^{a_i}}$ and $2^{\varphi(n)/2^k} \equiv 1 \pmod{\prod_{j=r+1}^k q_j^{b_j}}$, which means that $2^{\varphi(n)/2^k} \equiv +1 \pmod{\prod_{i=1}^r p_i^{a_i} \prod_{j=r+1}^k q_j^{b_j} = n}$, or $n \in (2^k+)$. $\square$

Note that if r in Theorem 2.5 was equal to zero, then we would have Theorem 2.3. So the last possibility is that r is equal to k , which we investigate in the following theorem.

Theorem 2.6. *Let $n = \prod p_i^{a_i} \prod q_j^{b_j}$ with $p_i \equiv 3 \pmod{8}$ for all i , and $q_j \equiv -1 \pmod{8}$ for all j , then*

$$n \notin (2^k+) \cup (2^k-).$$

Proof. Both $\varphi(p_i^{a_i})/2$ and $\varphi(q_j^{b_j})/2$ are odd, because $p_i \equiv q_j \equiv 3 \pmod{4}$, and so $\frac{p_i-1}{2} \equiv \frac{q_j-1}{2} \equiv 1 \pmod{2}$. So we have

$$2^{\varphi(n)/2^k} = (2^{\varphi(p_i^{a_i})/2})^{\frac{\prod_{l \neq i} \varphi(p_l^{a_l}) \prod \varphi(q_j^{b_j})}{2^{k-1}}} \equiv (-1)^{\frac{\prod_{l \neq i} \varphi(p_l^{a_l}) \prod \varphi(q_j^{b_j})}{2^{k-1}}} \equiv -1 \pmod{p_i^{a_i}},$$

because $\frac{\prod_{l \neq i} \varphi(p_l^{a_l}) \prod \varphi(q_j^{b_j})}{2^{k-1}}$ is odd. On the other hand,

$$2^{\varphi(n)/2^k} = (2^{\varphi(q_j^{b_j})/2})^{\frac{\prod_{l \neq j} \varphi(p_i^{a_i}) \prod \varphi(q_l^{b_l})}{2^{k-1}}} \equiv (+1)^{\frac{\prod_{l \neq j} \varphi(p_i^{a_i}) \prod \varphi(q_l^{b_l})}{2^{k-1}}} \equiv 1 \pmod{q_j^{b_j}}.$$

Thus, $2^{\varphi(n)/2^k} \equiv -1 \pmod{\prod p_i^{a_i}}$ and $2^{\varphi(n)/2^k} \equiv +1 \pmod{\prod q_j^{b_j}}$, which contradicts both $2^{\varphi(n)/2^k} \equiv +1 \pmod{n}$ and $2^{\varphi(n)/2^k} \equiv -1 \pmod{n}$; that is, $n \notin (2^k+) \cup (2^k-)$. $\square$

Note that Theorems 2.2, 2.3, 2.4, 2.5, and 2.6 cover all possible cases of n with $k \geq 2$ different prime divisors, and so Theorem 1 in the introduction is proved.

So the case when n has exactly k prime divisors is completely described. What can we say about the case when n has exactly $k-1$ prime divisors? Does n belong or not belong to the classes (2^k+) or (2^k-) in this case? Theorems 2.7, 2.8, 2.9, and 2.10 give a complete answer to this question.

From here on, we consider the case

$$n = \prod_{i=1}^{k-1} p_i^{a_i} \text{ where } k \geq 3.$$

In Theorems 2.7, and 2.8, we study the case in which n has at least one prime divisor congruent to 1 modulo 8. In Theorem 2.7, we assume that n has another prime divisor congruent to either 1 or -3 modulo 8, and in Theorem 2.8, we assume that all prime divisors but one, which is congruent to 1 as we already mentioned, are congruent to 1 or -3 modulo 8.

Theorem 2.7. *If n has at least two prime divisors, say p_1 and p_2 , with $p_1 \equiv 1 \pmod{8}$ and $p_2 \equiv 1$ or $-3 \pmod{8}$, then $n \in (2^k+)$.*

Proof. Let $n = p_1^{a_1} p_2^{a_2} A$, where $A = \prod_{i=3}^{k-1} p_i^{a_i}$ and $k \geq 3$. Since $p_1 \equiv 1 \pmod{8}$ and moreover, $\frac{\varphi(p_2^{a_2})}{4}$ is an integer (since $p_2 \equiv 1$ or $-3 \pmod{8}$ means that $p_2 - 1$ is divisible by 4), we have

$$2^{\frac{\varphi(n)}{2^k}} = (2^{\frac{\varphi(p_1^{a_1})}{2}})^{\frac{\varphi(p_2^{a_2})}{4} \frac{\varphi(A)}{2^{k-3}}} \equiv 1 \pmod{p_1^{a_1}}.$$

Since $\frac{\varphi(p_1^{a_1})}{4}$ is even ($8|p_1 - 1$), we have

$$2^{\frac{\varphi(n)}{2^k}} = \left(2^{\frac{\varphi(p_i^{a_i})}{2}}\right)^{\frac{\varphi(p_1^{a_1})}{4} \frac{\prod_{j=2, j \neq i}^{k-1} \varphi(p_j^{a_j})}{2^{k-3}}} \equiv 1 \pmod{p_i^{a_i}} \text{ for all } i = 2, \dots, k-1.$$

Thus, $2^{\frac{\varphi(n)}{2^k}} \equiv 1 \pmod{p_i^{a_i}}$ for all $i = 1, \dots, k-1$; therefore, $2^{\frac{\varphi(n)}{2^k}} \equiv +1 \pmod{n}$, in other words, $n \in (2^k+)$. $\square$

The following lemma give us an important property of the primes in the classes $(4+)$ and $(4-)$, which we need in the proof of Theorem 2.8.

Lemma 1. *A prime power $p^a \in (4+) \cup (4-)$ if and only if $p \equiv 1 \pmod{8}$.*

Proof. First, if $p^a \in (4+) \cup (4-)$, then $p \equiv 1 \pmod{8}$, is already proved in [1]. Conversely, if $p \equiv 1 \pmod{8}$, then $2^{\frac{p-1}{2}} \equiv 1 \pmod{p}$. So we can write

$$0 \equiv 2^{\frac{p-1}{2}} - 1 \equiv (2^{\frac{p-1}{4}} - 1)(2^{\frac{p-1}{4}} + 1) \pmod{p}.$$

Which means that $2^{\frac{p-1}{4}} \equiv +1$ or $-1 \pmod{p}$, that is $2^{\frac{p-1}{4}} = \pm 1 + Ap$ for some integer A . Thus

$$2^{\frac{\varphi(p^a)}{4}} = (2^{\frac{p-1}{4}})^{p^{a-1}} = (\pm 1 + Ap)^{p^{a-1}} = \pm 1 + Bp^a \text{ for some integer } B.$$

Then $p^a \in (4+) \cup (4-)$. $\square$

Theorem 2.8. *If $n = p_1^{a_1} \prod_{i=2}^{k-1} p_i^{a_i}$ with $p_1 \equiv 1 \pmod{8}$ and $p_i \equiv -1$ or $3 \pmod{8}$ for all $2 \leq i \leq k-1$, then*

$$\begin{aligned} n &\in (2^k+) && \text{if } p_1 \in (4+), \text{ and} \\ n &\notin (2^k+) \cup (2^k-) && \text{if } p_1 \in (4-). \end{aligned}$$

Proof. Let $n = p_1^{a_1} A$ where $A = \prod_{i=2}^{k-1} p_i^{a_i}$ with $p_i \equiv -1$ or $3 \pmod{8}$.

If $p_i \equiv -1 \pmod{8}$, then $2^{\frac{\varphi(p_i^{a_i})}{2}} \equiv 1 \pmod{p_i^{a_i}}$. Thus,

$$2^{\frac{\varphi(n)}{2^k}} = \left(2^{\frac{\varphi(p_i^{a_i})}{2}}\right)^{\frac{\varphi(p_1^{a_1})}{4} \frac{\prod_{j=2, j \neq i}^{k-1} \varphi(p_j^{a_j})}{2^{k-3}}} \equiv 1 \pmod{p_i^{a_i}}.$$

If $p_i \equiv 3 \pmod{8}$, then $2^{\frac{\varphi(p_i^{a_i})}{2}} \equiv -1 \pmod{p_i^{a_i}}$. Since $\frac{\varphi(p_1^{a_1})}{4}$ is even, we have

$$2^{\frac{\varphi(n)}{2^k}} = \left(2^{\frac{\varphi(p_i^{a_i})}{2}}\right)^{\frac{\varphi(p_1^{a_1})}{4} \frac{\prod_{j=2, j \neq i}^{k-1} \varphi(p_j^{a_j})}{2^{k-3}}} \equiv 1 \pmod{p_i^{a_i}}.$$

So $2^{\frac{\varphi(n)}{2^k}} \equiv 1 \pmod{p_i^{a_i}}$ for all $i = 2, \dots, k-1$. Therefore,

$$2^{\frac{\varphi(n)}{2^k}} \equiv 1 \pmod{A}.$$

Since $\frac{\varphi(p_i^{a_i})}{2}$ is odd for all $i = 2, \dots, k-1$, only $\varphi(p_1^{a_1})$ can be divisible by 4. Since $p^a \in (4+) \cup (4-)$ if and only if $p \equiv 1 \pmod{8}$, as we proved in Lemma 1, we have $p_1 \in (4+)$ or $\in (4-)$.

If $p_1 \in (4+)$, then $2^{\frac{\varphi(p_1^{a_1})}{4}} \equiv +1 \pmod{p_1^{a_1}}$. Therefore,

$$2^{\frac{\varphi(n)}{2^k}} = \left(2^{\frac{\varphi(p_1^{a_1})}{4}}\right)^{\frac{\varphi(A)}{2^{k-2}}} \equiv (+1)^{\frac{\varphi(A)}{2^{k-2}}} \equiv 1 \pmod{p_1^{a_1}}.$$

So now we have $2^{\frac{\varphi(n)}{2^k}} \equiv 1 \pmod{p_1^{a_1}}$ and $2^{\frac{\varphi(n)}{2^k}} \equiv 1 \pmod{A}$, which means that $2^{\frac{\varphi(n)}{2^k}} \equiv +1 \pmod{p_1^{a_1}A = n}$, or $n \in (2^k+)$.

If $p_1 \in (4-)$, then $2^{\frac{\varphi(p_1^{a_1})}{4}} \equiv -1 \pmod{p_1^{a_1}}$. Therefore,

$$2^{\frac{\varphi(n)}{2^k}} = \left(2^{\frac{\varphi(p_1^{a_1})}{4}}\right)^{\frac{\varphi(A)}{2^{k-2}}} \equiv (-1)^{\frac{\varphi(A)}{2^{k-2}}} \equiv -1 \pmod{p_1^{a_1}},$$

because $\frac{\varphi(A)}{2^{k-2}}$ is odd. So we have $2^{\frac{\varphi(n)}{2^k}} \equiv -1 \pmod{p_1^{a_1}}$ and $2^{\frac{\varphi(n)}{2^k}} \equiv +1 \pmod{A}$. So $n \notin (2^k+) \cup (2^k-)$. Indeed, from $n \in (2^k+) \cup (2^k-)$, it follows that $2^{\frac{\varphi(n)}{2^k}} \equiv \pm 1 \pmod{p_1^{a_1}}$ and $2^{\frac{\varphi(n)}{2^k}} \equiv \pm 1 \pmod{A}$, which contradicts what we have. $\square$

Remark: Since $p \equiv 1 \pmod{8}$ is equivalent to $p \in (4+) \cup (4-)$, we can reformulate Theorems 2.7 and 2.8 as follows.

- If n has at least one prime divisor $p \in (4+)$, then $n \in (2^k+)$.
- If n has at least two prime divisors p_1 and p_2 with $p_1 \in (4-)$ and $p_2 \equiv 1$ or $-3 \pmod{8}$, then $n \in (2^k+)$.
- If n has one prime divisor p_1 with $p_1 \in (4-)$ and all the other prime divisors $p_i \equiv -1$ or $3 \pmod{8}$, then $n \notin (2^k+) \cup (2^k-)$.

finally, in Theorems 2.9 and 2.10 we consider the case in which n has no prime divisor congruent to 1 modulo 8.

Theorem 2.9. *If $n = \prod_{i=1}^{k-1} p_i^{a_i}$ with $p_i \equiv -3 \pmod{8}$ for all i , then*

$$\begin{aligned} n &\in (2^k+) && \text{if } k > 3, \text{ and} \\ n &\in (2^k-) && \text{if } k = 3. \end{aligned}$$

Proof. Since $p_i \equiv -3 \pmod{8}$, it follows that $2^{\frac{\varphi(p_i^{a_i})}{2}} \equiv -1 \pmod{p_i^{a_i}}$, $\frac{\varphi(p_i^{a_i})}{2}$ is even, but $\frac{\varphi(p_i^{a_i})}{4}$ is odd. Hence,

$$2^{\frac{\varphi(n)}{2^k}} = (2^{\frac{\varphi(p_i^{a_i})}{2}})^{\frac{\varphi(p_1^{a_1})}{4} \frac{\prod_{j=1, j \neq i, l}^{k-1} \varphi(p_j^{a_j})}{2^{k-3}}} \equiv (-1)^{\frac{\varphi(p_l^{a_l})}{4} \frac{\prod_{j=1, j \neq i, l}^{k-1} \varphi(p_j^{a_j})}{2^{k-3}}} \pmod{p_i^{a_i}}.$$

If $k > 3$, then $\frac{\prod_{j=1, j \neq i, l}^{k-1} \varphi(p_j^{a_j})}{2^{k-3}}$ is even. Thus, $2^{\frac{\varphi(n)}{2^k}} \equiv +1 \pmod{p_i^{a_i}}$ for all i , and then $2^{\frac{\varphi(n)}{2^k}} \equiv +1 \pmod{n}$, or $n \in (2^k+)$.

If $k = 3$, then $n = p_1^{a_1} p_2^{a_2}$. Therefore,

$$2^{\frac{\varphi(n)}{2^3}} = (2^{\frac{\varphi(p_1^{a_1})}{2}})^{\frac{\varphi(p_2^{a_2})}{4}} \equiv -1 \pmod{p_1^{a_1}}, \text{ and } 2^{\frac{\varphi(n)}{2^3}} = (2^{\frac{\varphi(p_2^{a_2})}{2}})^{\frac{\varphi(p_1^{a_1})}{4}} \equiv -1 \pmod{p_2^{a_2}}.$$

Thus, $2^{\frac{\varphi(n)}{2^3}} \equiv -1 \pmod{p_1^{a_1} p_2^{a_2} = n}$, or $n \in (2^k-)$. $\square$

Theorem 2.10. *If $n = \prod_{i=1}^r p_i^{a_i} \prod_{j=r+1}^{k-1} q_j^{b_j}$ with $0 \leq r < k-1$, $p_i \equiv -3 \pmod{8}$ for all i , and $q_j \equiv -1$ or $3 \pmod{8}$ for all j , then*

$$\begin{aligned} n &\notin (2^k+) \cup (2^k-) && \text{if } 0 \leq r \leq 2, \text{ and} \\ n &\in (2^k+) && \text{if } 2 < r < k-1. \end{aligned}$$

Proof. Now we have that $\varphi(p_i^{a_i})/2$ is even and $\varphi(p_i^{a_i})/4$ is odd. Indeed, $p_i \equiv -3 \pmod{8}$ means that $p_i = 5 + 8s_i$ for some integer s_i , it follows that $\frac{p_i-1}{2} = 2 + 4s_i$ is even and $\frac{p_i-1}{4} = 1 + 2s_i$ is odd. Therefore, $\frac{\varphi(p_i^{a_i})}{2} = \frac{p_i-1}{2} p_i^{a_i-1}$ is even and $\frac{\varphi(p_i^{a_i})}{4} = \frac{p_i-1}{4} p_i^{a_i-1}$ is odd.

Moreover, $\varphi(q_j^{b_j})/2$ is odd. Indeed, $q_j \equiv -1$ or $3 \pmod{8}$ means that $q_j \equiv 3 \pmod{4}$ for all j , so $\frac{q_j-1}{2}$ is odd, and therefore $\frac{q_j-1}{2} q_j^{b_j-1} = \varphi(q_j^{b_j})/2$ is odd.

If $r = 0$, then $n = \prod_{j=1}^{k-1} q_j^{b_j}$ and $\varphi(n) = \prod_{j=1}^{k-1} \varphi(q_j^{b_j})$. Since $\varphi(q_j^{b_j})/2$ is odd for all j , it follows that $\frac{\prod_{j=1}^{k-1} \varphi(q_j^{b_j})}{2^{k-1}}$ is odd. Thus, $\varphi(n)$ is not divisible by 2^k , which implies that $n \notin (2^k+) \cup (2^k-)$.

If $r = 1$, then $n = p_1^{a_1} \prod_{j=2}^{k-1} q_j^{b_j}$ and $\frac{\varphi(n)}{2^k} = \frac{\varphi(p_1^{a_1})}{4} \frac{\prod_{j=2}^{k-1} \varphi(q_j^{b_j})}{2^{k-2}}$. So

$$2^{\frac{\varphi(n)}{2^k}} = (2^{\frac{\varphi(q_j^{b_j})}{2}})^{\frac{\varphi(p_1^{a_1})}{4} \frac{\prod_{l=2, l \neq j}^{k-1} \varphi(q_l^{b_l})}{2^{k-3}}} \equiv (\pm 1)^{\frac{\varphi(p_1^{a_1})}{4} \frac{\prod_{l=2, l \neq j}^{k-1} \varphi(q_l^{b_l})}{2^{k-3}}} \equiv \pm 1 \pmod{q_j^{b_j}},$$

because $\frac{\varphi(p_1^{a_1})}{4} \frac{\prod_{l=2, l \neq j}^{k-1} \varphi(q_l^{b_l})}{2^{k-3}}$ is odd. On the other hand,

$$2^{\frac{\varphi(n)}{2^k}} = (2^{\frac{\varphi(p_1^{a_1})}{4}})^{\frac{\prod_{j=2}^{k-1} \varphi(q_j^{b_j})}{2^{k-2}}} \not\equiv +1 \text{ or } -1 \pmod{p_1^{a_1}}.$$

Indeed, $p_1^{a_1} \notin (4+) \cup (4-) \cup (2+)$, because $p_1 \equiv -3 \pmod{8}$; therefore, $2^{\frac{\varphi(p_1^{a_1})}{4}} \not\equiv \pm 1 \pmod{p_1^{a_1}}$, and $2^{\frac{\varphi(p_1^{a_1})}{2}} \not\equiv +1 \pmod{p_1^{a_1}}$. Hence,

$2^{\frac{\varphi(n)}{2^k}} = (2^{\frac{\varphi(p_1^{a_1})}{4}})^{\frac{\prod_{j=2}^{k-1} \varphi(q_j^{b_j})}{2^{k-2}}} \equiv +1 \pmod{p_1^{a_1}}$ only if $\frac{\prod_{j=2}^{k-1} \varphi(q_j^{b_j})}{2^{k-2}}$ is divisible by 4.

But $\frac{\prod_{j=2}^{k-1} \varphi(q_j^{b_j})}{2^{k-2}}$ is odd, so $2^{\frac{\varphi(n)}{2^k}} \not\equiv +1 \pmod{p_1^{a_1}}$.

Since $2^{\frac{\varphi(p_1^{a_1})}{2}} \equiv -1 \pmod{p_1^{a_1}}$, we have $2^{\frac{\varphi(n)}{2^k}} \equiv -1$ only if $\frac{\prod_{j=2}^{k-1} \varphi(q_j^{b_j})}{2^{k-1}}$ is an odd integer. But $\frac{\prod_{j=2}^{k-1} \varphi(q_j^{b_j})}{2^{k-2}}$ is odd, which implies that $2^{k-1} \nmid \prod_{j=2}^{k-1} \varphi(q_j^{b_j})$; therefore, $2^{\frac{\varphi(n)}{2^k}} \not\equiv -1 \pmod{p_1^{a_1}}$.

So we have $2^{\frac{\varphi(n)}{2^k}} \equiv \pm 1 \pmod{q_j^{b_j}}$ and $2^{\frac{\varphi(n)}{2^k}} \not\equiv \pm 1 \pmod{p_1^{a_1}}$, which implies that $2^{\frac{\varphi(n)}{2^k}} \not\equiv \pm 1 \pmod{n}$, i.e. $n \notin (2^k+) \cup (2^k-)$.

If $r = 2$, then $n = p_1^{a_1} p_2^{a_2} \prod_{j=3}^{k-1} q_j^{b_j}$ and $\frac{\varphi(n)}{2^k} = \frac{\varphi(p_1^{a_1})\varphi(p_2^{a_2})}{8} (\frac{\prod_{j=3}^{k-1} \varphi(q_j^{b_j})}{2^{k-3}})$. So

$$2^{\frac{\varphi(n)}{2^k}} = (2^{\frac{\varphi(p_1^{a_1})}{2}})^{\frac{\varphi(p_2^{a_2})}{4} \frac{\prod_{j=3}^{k-1} \varphi(q_j^{b_j})}{2^{k-3}}} \equiv (-1)^{\frac{\varphi(p_2^{a_2})}{4} \frac{\prod_{j=3}^{k-1} \varphi(q_j^{b_j})}{2^{k-3}}} \equiv -1 \pmod{p_1^{a_1}},$$

because $\frac{\varphi(p_2^{a_2})}{4} \frac{\prod_{j=3}^{k-1} \varphi(q_j^{b_j})}{2^{k-3}}$ is odd. Also,

$$2^{\frac{\varphi(n)}{2^k}} = (2^{\frac{\varphi(p_2^{a_2})}{2}})^{\frac{\varphi(p_1^{a_1})}{4} \frac{\prod_{j=3}^{k-1} \varphi(q_j^{b_j})}{2^{k-3}}} \equiv (-1)^{\frac{\varphi(p_1^{a_1})}{4} \frac{\prod_{j=3}^{k-1} \varphi(q_j^{b_j})}{2^{k-3}}} \equiv -1 \pmod{p_2^{a_2}}.$$

On the other hand,

$$\begin{aligned} 2^{\frac{\varphi(n)}{2^k}} &= (2^{\frac{\varphi(q_j^{b_j})}{2}})^{\frac{\varphi(p_1^{a_1})\varphi(p_2^{a_2})}{8} \frac{\prod_{l=3, l \neq j}^{k-1} \varphi(q_l^{b_l})}{2^{k-4}}} \equiv (\pm 1)^{\frac{\varphi(p_1^{a_1})\varphi(p_2^{a_2})}{8} \frac{\prod_{l=3, l \neq j}^{k-1} \varphi(q_l^{b_l})}{2^{k-4}}} \\ &\equiv +1 \pmod{q_j^{b_j}}, \text{ because } \frac{\varphi(p_1^{a_1})\varphi(p_2^{a_2})}{8} \text{ is even.} \end{aligned}$$

Therefore, $2^{\frac{\varphi(n)}{2^k}} \equiv +1 \pmod{q_j^{b_j}}$ and $2^{\frac{\varphi(n)}{2^k}} \equiv -1 \pmod{p_i^{a_i}}$ which contradicts both $2^{\frac{\varphi(n)}{2^k}} \equiv +1$ and $-1 \pmod{n}$.

Thus, $n \notin (2^k+) \cup (2^k-)$.

If $2 < r < k - 1$, then

$$\begin{aligned} 2^{\frac{\varphi(n)}{2^k}} &= (2^{\frac{\varphi(p_i^{a_i})}{2}})^{\frac{\prod_{l=1, l \neq i}^r \varphi(p_l^{a_l})}{2^r} \frac{\prod_{j=r+1}^{k-1} \varphi(q_j^{b_j})}{2^{k-r-1}}} \equiv (-1)^{\frac{\prod_{l=1, l \neq i}^r \varphi(p_l^{a_l})}{2^r} \frac{\prod_{j=r+1}^{k-1} \varphi(q_j^{b_j})}{2^{k-r-1}}} \\ &\equiv +1 \pmod{p_i^{a_i}}, \text{ because } \frac{\prod_{l=1, l \neq i}^r \varphi(p_l^{a_l})}{2^r} \text{ is even.} \end{aligned}$$

$$\begin{aligned} \text{Also, } 2^{\frac{\varphi(n)}{2^k}} &= (2^{\frac{\varphi(q_j^{b_j})}{2}})^{\frac{\prod_{i=1}^r \varphi(p_i^{a_i})}{2^{r+1}} \frac{\prod_{l=r+1, l \neq j}^{k-1} \varphi(q_l^{b_l})}{2^{k-r-2}}} \equiv (\pm 1)^{\frac{\prod_{i=1}^r \varphi(p_i^{a_i})}{2^{r+1}} \frac{\prod_{l=r+1, l \neq j}^{k-1} \varphi(q_l^{b_l})}{2^{k-r-2}}} \\ &\equiv +1 \pmod{q_j^{b_j}}, \text{ because } \frac{\prod_{i=1}^r \varphi(p_i^{a_i})}{2^{r+1}} \text{ is even.} \end{aligned}$$

So $2^{\frac{\varphi(n)}{2^k}} \equiv +1 \pmod{p_i^{a_i}}$ and $2^{\frac{\varphi(n)}{2^k}} \equiv +1 \pmod{q_j^{b_j}}$ for all i and j . Thus, $2^{\frac{\varphi(n)}{2^k}} \equiv +1 \pmod{n}$, or $n \in (2^k+)$. $\square$

We can easily see that Theorems 2.7, 2.8, 2.9, and 2.10 cover all possible cases of n with $k-1 \geq 2$ different prime divisors. Therefore, the proof of Theorem 2 in the introduction is complete.

Remark: Using our theorems, it is easy to check whether $n \in (2^k+)$, $n \in (2^k-)$, or $n \notin (2^k+) \cup (2^k-)$, in case when n has $k-1$ or more distinct prime divisors. If n has less than $k-1$ distinct prime divisors, then we can easily see that $n \notin (2^k+) \cup (2^k-)$, if $n = \prod_{i=1}^r p_i^{a_i}$, with $0 < r < k-1$ and $p_i \equiv -1$ or $3 \pmod{8}$ for all i , because in this case $2^k \nmid \varphi(n)$.

3 The classes $(2\pm)$, $(4\pm)$ and $(8\pm)$

The classes $(2\pm)$ or the case $k=1$ is completely described in section 2. Now let us study the case $k=2$ or the classes $(4\pm)$.

The following properties are proved in [1], and we will see that, they can be obtained as special cases of the theorems in section 2.

P1. Every odd number, divisible by more than two primes, belongs to the class $(4+)$.

The following 5 properties describe the case in which n is divisible by two different primes p, q .

P2. If $p \equiv +1 \pmod{8}$, then $n \in (4+)$.

P3. If $p \equiv q \equiv -1 \pmod{8}$ or $p \equiv q \equiv -3 \pmod{8}$, then $n \in (4+)$.

P4. If $p \equiv q \equiv +3 \pmod{8}$, then $n \in (4-)$.

P5. If $p \equiv +3 \pmod{8}, q \equiv -1$ or $-3 \pmod{8}$, then $n \notin (4+) \cup (4-)$.

P6. If $p \equiv -3 \pmod{8}, q \equiv -1 \pmod{8}$, then $n \notin (4+) \cup (4-)$.

The following property considers the case in which n has only one prime divisor.

P7. If $n = p^a$ belongs to the class $(4+) \cup (4-)$, then $p \equiv 1 \pmod{8}$.

The following two properties consider every odd number n .

P8. If $n \equiv 5 \pmod{8}$, then n does not belong to the class $(4-)$.

P9. If $n \equiv 7 \pmod{8}$, then n does not belong to the class $(4-)$.

- It is clear that if $k = 2$, then P1, P2, P3, and P4 are special cases from theorems 2.1, 2.2, 2.3, and 2.4 respectively.
- In P5 we have two cases:
 - Case 1: If $p \equiv +3 \pmod{8}, q \equiv -1 \pmod{8}$, which is exactly theorem 2.6 with $k = 2$.
 - Case 2: If $p \equiv +3 \pmod{8}, q \equiv -3 \pmod{8}$, which is exactly the case in theorem 2.5 with $k = 2$ and $r = 1$.
- The case in P6 is the case in theorem 2.5 with $k = 2$ and $r = 1$.
- In P7, we have just one direction and Lemma 1 complete the second direction.
- In P8 and P9, n has three cases. Case 1: If n has more than two distinct prime divisors, then by theorem 2.2, n belongs to $(4+)$ not to $(4-)$.
 - Case 2: If n has two prime divisors, then by theorem 2.4, n belongs to $(4-)$ only if all the prime divisors of n are congruent to $3 \pmod{8}$. Since $3^2 \equiv 1 \pmod{8}$, we have $n \in (4-)$ only if $n \equiv 3$ or $1 \pmod{8}$. So $n \notin (4-)$ if $n \equiv 5$ or $7 \pmod{8}$.
 - Case 3: If n has only one prime divisor, then by Lemma 1 $n \notin (4+) \cup (4-)$.

The classes $(8\pm)$:

If $k = 3$, theorem 2.1 state the following:

Every odd number, divisible by more than three primes, belongs to the class $(8+)$.

So now consider the case $n = p^a q^b s^c, a > 0, b > 0, c > 0$. In [2], Arnold denoted by I, II, III, IV the sets of the primes, congruent to 1, 3, 5 and to 7 mod 8 respectively, and proved the following properties:

- P1. The triple product n belongs to the class $(8-)$, whenever the primes (p, q, s) do all belong to the kind II.
- P2. The triple product n belongs to the class $(8+)$, provided that the three primes p, q and s do belong to one of the following 7 triples of kinds:
 (I, I, X), (I, III, X), (I, II, II), (I, II, IV), (I, IV, IV), (IV, IV, IV),
 (III, III, X).

P3. No triple product n belongs to the class $(8+)$ or $(8-)$, provided that the primes (p, q, s) kinds triple is one of the 5 triples:

$(\text{III}, \text{II}, \text{II}), (\text{III}, \text{II}, \text{IV}), (\text{III}, \text{IV}, \text{IV}), (\text{II}, \text{IV}, \text{II}), (\text{II}, \text{IV}, \text{IV})$.

Note that the triples in P1, P2, and P3 cover (taking the permutations into account) all the 64 possible ordered lists of the 3 kinds of p , q , and s . To verify it, see [2].

Now consider the case if n has exactly $k - 1 = 2$ distinct prime divisors. Let $n = p^a q^b$, $a > 0, b > 0$. In [2] Arnold proved the following properties:

P4. If $p \in \text{III}$ and $q \in \text{III}$, then $n \in (8-)$.

P5. If $p \in \text{I}$ and $q \in \text{I} \cup \text{III}$, then $n \in (8+)$.

P6. If $p \in (\text{I} \cap (4+))$ and $q \in (\text{II} \cup \text{IV})$, then $n \in (8+)$.

P7. If $p \in (\text{I} \cap (4-))$ and $q \in (\text{II} \cup \text{IV})$, then $n \notin (8+) \cup (8-)$.

P8. If $p \in (\text{II} \cup \text{IV})$ and $q \in (\text{II} \cup \text{IV})$, then $n \notin (8+) \cup (8-)$.

P9. If $p \in (4+) \cup (4-)$ and $q \in (4+) \cup (4-)$, then $pq \in (8+)$.

We will see now that these properties are special cases of the theorems in section 2.

- It is clear that P1 is exactly theorem 2.4 with $k = 3$.
- In P2, the first 5 triples $(\text{I}, \text{I}, \text{X}), (\text{I}, \text{III}, \text{X}), (\text{I}, \text{II}, \text{II}), (\text{I}, \text{II}, \text{IV}), (\text{I}, \text{IV}, \text{IV})$ have at least one prime divisor of kind I, that is they satisfy the condition of theorem 2.2. So they are special cases of it.

The triple $(\text{IV}, \text{IV}, \text{IV})$ is exactly theorem 2.3 with $k = 3$.

The triple $(\text{III}, \text{III}, \text{X})$ has 3 cases. Case 1: the triple $(\text{III}, \text{III}, \text{I})$ clearly belongs to theorem 2.2.

Case 2: the triples $(\text{III}, \text{III}, \text{II})$ and $(\text{III}, \text{III}, \text{IV})$ have two prime divisors of kind III and only one divisor of kind II or IV, so they belong to theorem 2.5 with $k = 3, r = 1 < k - 1$.

Case 3: the triple $(\text{III}, \text{III}, \text{III})$ is exactly theorem 2.3 with $k = 3$.

- In P3, the first 3 triples $(\text{III}, \text{II}, \text{II}), (\text{III}, \text{II}, \text{IV}), (\text{III}, \text{IV}, \text{IV})$ have one prime divisor from kind III and the others from kind II or IV, which means that they are special cases from theorem 2.5 with $k = 3, r = k - 1 = 2$.

The triples $(\text{II}, \text{IV}, \text{II}), (\text{II}, \text{IV}, \text{IV})$ have only prime divisors from kinds II or IV, so they belong to theorem 2.6 with $k = 3$.

- It is clear that the properties P4-P9 can be obtained from Theorems 2.7-2.10 and Lemma 1.

References

- [1] ARNOLD, V. I., *Fermat-Euler Dynamical Systems and the Statistics of Arithmetics of Geometric Progressions*, Funktsional. Anal. i Prilozhen. **37** (2003), no. 1, 1–18, 95; translation in Funct. Anal. Appl. **37** (2003), no. 1, 1–15.
- [2] ARNOLD, V. I., *Ergodic and Arithmetical properties of geometrical progression's dynamics and its orbits*, Mosc. Math. J. **5** (2005), no. 1, 5–22.
- [3] SILVERMAN, JOSEPH H., *A Friendly Introduction to Number Theory*, Prentice Hall, **1996**.